

## **Synthesis of novel asymmetrical 1, 4-dihydropyridine derivatives as anti tubercular agents and evaluation of their calcium channel blocking activity**

Ramin Miri<sup>a</sup>, Hossein Sadegh pour<sup>a</sup>, Katayoun Javidnia<sup>a</sup>, Omid reza Firouzi<sup>b</sup>, Ali reza Nikkhoo<sup>c</sup>

<sup>a</sup>Department of medicinal chemistry, School of Pharmacy, Shiraz University of Medical Science

<sup>b</sup>Medicinal & Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

One third of world population is infected with latent tuberculosis (TB) infection. Based on this fact, many people are at risk to develop active TB. In 2011, tuberculosis (TB) remained the second cause of death from infectious disease worldwide. Every second one person is infected and every fifteen second someone dies with TB. Therefore, the search for new antimycobacterials is the subject of numerous recent studies.

Previous studies showed that 1, 4-dihydropyridine containing alkyl ester and diethyl carbamoyl could effectively overcome MTB. Thus, the asymmetrical analogues (6a-m) of 1, 4-dihydropyridines were synthesized by a modified Hantzsch reaction using a procedure reported by Iwanami. Reaction of acetoacetates with ammonium acetate afforded the corresponding 3-aminocrotonates. Subsequent reaction of 3-aminocrotonates with aldehydes and N, N-dimethyl acetoacetamide afforded final 1, 4-dihydropyridines (**6a-m**). The obtained compounds were purified by preparative thin layer chromatography and recrystallized from ethyl acetate-petroleum ether. They were identified by <sup>1</sup>HNMR, mass spectroscopy and IR.

These compounds have essential requirements for calcium channel blocker activity as dihydropyridine class. Therefore, these compounds were evaluated as calcium-channel antagonists by using high potassium ion concentration in guinea-pig ileal longitudinal smooth muscle. They all exhibited less calcium channel antagonist activity (IC<sub>50</sub>=10<sup>-5</sup> to

$10^{-6}$ ) relative to the reference drug Nifedipine ( $6.31 \times 10^{-8}$ ). Docking studies offer the binding pattern of compounds to the InhA from MTB and L-type  $\text{Ca}^{2+}$  channel. The result of pharmacological assay was used in quantitative structure-activity relationship studies utilizing multiple linear regression analysis. QSAR study indicates that the activity is related to the electronic and topological parameters. These compounds also evaluated as antimicrobial agents and they were inactive.

**Keywords:** Tuberculosis, 1, 4-dihydropyridines, calcium channel blockers, docking quantitative structure-activity relationship studie

## **Synthesis of novel asymmetrical 1, 4-dihydropyridine derivatives as anti tubercular agents and evaluation of their calcium channel blocking activity**

Ramin Miri<sup>a</sup>, Hossein Sadegh pour<sup>a</sup>, Katayoun Javidnia<sup>a</sup>, Omid reza Firouzi<sup>b</sup>, Ali reza Nikkhoo<sup>c</sup>

<sup>a</sup>Department of medicinal chemistry, School of Pharmacy, Shiraz University of Medical Science

<sup>b</sup>Medicinal & Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

One third of world population is infected with latent tuberculosis (TB) infection. Based on this fact, many people are at risk to develop active TB. In 2011, tuberculosis (TB) remained the second cause of death from infectious disease worldwide. Every second one person is infected and every fifteen second someone dies with TB. Therefore, the search for new antimycobacterials is the subject of numerous recent studies.

Previous studies showed that 1, 4-dihydropyridine containing alkyl ester and diethyl carbamoyl could effectively overcome MTB. Thus, the asymmetrical analogues (6a-m) of 1, 4-dihydropyridines were synthesized by a modified Hantzsch reaction using a procedure reported by Iwanami. Reaction of acetoacetates with ammonium acetate afforded the corresponding 3-aminocrotonates. Subsequent reaction of 3-aminocrotonates with aldehydes and N, N-dimethyl acetoacetamide afforded final 1, 4-dihydropyridines (**6a-m**). The obtained compounds were purified by preparative thin layer chromatography and recrystallized from ethyl acetate-petroleum ether. They were identified by <sup>1</sup>HNMR, mass spectroscopy and IR.

These compounds have essential requirements for calcium channel blocker activity as dihydropyridine class. Therefore, these compounds were evaluated as calcium-channel antagonists by using high potassium ion concentration in guinea-pig ileal longitudinal smooth muscle. They all exhibited less calcium channel antagonist activity ( $IC_{50}=10^{-5}$  to

$10^{-6}$ ) relative to the reference drug Nifedipine ( $6.31 \times 10^{-8}$ ). Docking studies offer the binding pattern of compounds to the InhA from MTB and L-type  $\text{Ca}^{2+}$  channel. The result of pharmacological assay was used in quantitative structure-activity relationship studies utilizing multiple linear regression analysis. QSAR study indicates that the activity is related to the electronic and topological parameters. These compounds also evaluated as antimicrobial agents and they were inactive.

**Keywords:** Tuberculosis, 1, 4-dihydropyridines, calcium channel blockers, docking quantitative structure-activity relationship studie

## **Synthesis of novel asymmetrical 1, 4-dihydropyridine derivatives as anti tubercular agents and evaluation of their calcium channel blocking activity**

Ramin Miri<sup>a</sup>, Hossein Sadegh pour<sup>a</sup>, Katayoun Javidnia<sup>a</sup>, Omid reza Firouzi<sup>b</sup>, Ali reza Nikkhoo<sup>c</sup>

<sup>a</sup>Department of medicinal chemistry, School of Pharmacy, Shiraz University of Medical Science

<sup>b</sup>Medicinal & Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

One third of world population is infected with latent tuberculosis (TB) infection. Based on this fact, many people are at risk to develop active TB. In 2011, tuberculosis (TB) remained the second cause of death from infectious disease worldwide. Every second one person is infected and every fifteen second someone dies with TB. Therefore, the search for new antimycobacterials is the subject of numerous recent studies.

Previous studies showed that 1, 4-dihydropyridine containing alkyl ester and diethyl carbamoyl could effectively overcome MTB. Thus, the asymmetrical analogues (6a-m) of 1, 4-dihydropyridines were synthesized by a modified Hantzsch reaction using a procedure reported by Iwanami. Reaction of acetoacetates with ammonium acetate afforded the corresponding 3-aminocrotonates. Subsequent reaction of 3-aminocrotonates with aldehydes and N, N-dimethyl acetoacetamide afforded final 1, 4-dihydropyridines (**6a-m**). The obtained compounds were purified by preparative thin layer chromatography and recrystallized from ethyl acetate-petroleum ether. They were identified by <sup>1</sup>HNMR, mass spectroscopy and IR.

These compounds have essential requirements for calcium channel blocker activity as dihydropyridine class. Therefore, these compounds were evaluated as calcium-channel antagonists by using high potassium ion concentration in guinea-pig ileal longitudinal smooth muscle. They all exhibited less calcium channel antagonist activity ( $IC_{50}=10^{-5}$  to

$10^{-6}$ ) relative to the reference drug Nifedipine ( $6.31 \times 10^{-8}$ ). Docking studies offer the binding pattern of compounds to the InhA from MTB and L-type  $\text{Ca}^{2+}$  channel. The result of pharmacological assay was used in quantitative structure-activity relationship studies utilizing multiple linear regression analysis. QSAR study indicates that the activity is related to the electronic and topological parameters. These compounds also evaluated as antimicrobial agents and they were inactive.

**Keywords:** Tuberculosis, 1, 4-dihydropyridines, calcium channel blockers, docking quantitative structure-activity relationship studie

## **Synthesis of novel asymmetrical 1, 4-dihydropyridine derivatives as anti tubercular agents and evaluation of their calcium channel blocking activity**

Ramin Miri<sup>a</sup>, Hossein Sadegh pour<sup>a</sup>, Katayoun Javidnia<sup>a</sup>, Omid reza Firouzi<sup>b</sup>, Ali reza Nikkhoo<sup>c</sup>

<sup>a</sup>Department of medicinal chemistry, School of Pharmacy, Shiraz University of Medical Science

<sup>b</sup>Medicinal & Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

One third of world population is infected with latent tuberculosis (TB) infection. Based on this fact, many people are at risk to develop active TB. In 2011, tuberculosis (TB) remained the second cause of death from infectious disease worldwide. Every second one person is infected and every fifteen second someone dies with TB. Therefore, the search for new antimycobacterials is the subject of numerous recent studies.

Previous studies showed that 1, 4-dihydropyridine containing alkyl ester and diethyl carbamoyl could effectively overcome MTB. Thus, the asymmetrical analogues (6a-m) of 1, 4-dihydropyridines were synthesized by a modified Hantzsch reaction using a procedure reported by Iwanami. Reaction of acetoacetates with ammonium acetate afforded the corresponding 3-aminocrotonates. Subsequent reaction of 3-aminocrotonates with aldehydes and N, N-dimethyl acetoacetamide afforded final 1, 4-dihydropyridines (**6a-m**). The obtained compounds were purified by preparative thin layer chromatography and recrystallized from ethyl acetate-petroleum ether. They were identified by <sup>1</sup>HNMR, mass spectroscopy and IR.

These compounds have essential requirements for calcium channel blocker activity as dihydropyridine class. Therefore, these compounds were evaluated as calcium-channel antagonists by using high potassium ion concentration in guinea-pig ileal longitudinal smooth muscle. They all exhibited less calcium channel antagonist activity ( $IC_{50}=10^{-5}$  to

$10^{-6}$ ) relative to the reference drug Nifedipine ( $6.31 \times 10^{-8}$ ). Docking studies offer the binding pattern of compounds to the InhA from MTB and L-type  $\text{Ca}^{2+}$  channel. The result of pharmacological assay was used in quantitative structure-activity relationship studies utilizing multiple linear regression analysis. QSAR study indicates that the activity is related to the electronic and topological parameters. These compounds also evaluated as antimicrobial agents and they were inactive.

**Keywords:** Tuberculosis, 1, 4-dihydropyridines, calcium channel blockers, docking quantitative structure-activity relationship studie



## **Synthesis of novel asymmetrical 1, 4-dihydropyridine derivatives as anti tubercular agents and evaluation of their calcium channel blocking activity**

Ramin Miri<sup>a</sup>, Hossein Sadegh pour<sup>a</sup>, Katayoun Javidnia<sup>a</sup>, Omid reza Firouzi<sup>b</sup>, Ali reza Nikkhoo<sup>c</sup>

<sup>a</sup>Department of medicinal chemistry, School of Pharmacy, Shiraz University of Medical Science

<sup>b</sup>Medicinal & Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

One third of world population is infected with latent tuberculosis (TB) infection. Based on this fact, many people are at risk to develop active TB. In 2011, tuberculosis (TB) remained the second cause of death from infectious disease worldwide. Every second one person is infected and every fifteen second someone dies with TB. Therefore, the search for new antimycobacterials is the subject of numerous recent studies.

Previous studies showed that 1, 4-dihydropyridine containing alkyl ester and diethyl carbamoyl could effectively overcome MTB. Thus, the asymmetrical analogues (6a-m) of 1, 4-dihydropyridines were synthesized by a modified Hantzsch reaction using a procedure reported by Iwanami. Reaction of acetoacetates with ammonium acetate afforded the corresponding 3-aminocrotonates. Subsequent reaction of 3-aminocrotonates with aldehydes and N, N-dimethyl acetoacetamide afforded final 1, 4-dihydropyridines (**6a-m**). The obtained compounds were purified by preparative thin layer chromatography and recrystallized from ethyl acetate-petroleum ether. They were identified by <sup>1</sup>HNMR, mass spectroscopy and IR.

These compounds have essential requirements for calcium channel blocker activity as dihydropyridine class. Therefore, these compounds were evaluated as calcium-channel antagonists by using high potassium ion concentration in guinea-pig ileal longitudinal smooth muscle. They all exhibited less calcium channel antagonist activity ( $IC_{50}=10^{-5}$  to

$10^{-6}$ ) relative to the reference drug Nifedipine ( $6.31 \times 10^{-8}$ ). Docking studies offer the binding pattern of compounds to the InhA from MTB and L-type  $\text{Ca}^{2+}$  channel. The result of pharmacological assay was used in quantitative structure-activity relationship studies utilizing multiple linear regression analysis. QSAR study indicates that the activity is related to the electronic and topological parameters. These compounds also evaluated as antimicrobial agents and they were inactive.

**Keywords:** Tuberculosis, 1, 4-dihydropyridines, calcium channel blockers, docking quantitative structure-activity relationship studie

## **Synthesis of novel asymmetrical 1, 4-dihydropyridine derivatives as anti tubercular agents and evaluation of their calcium channel blocking activity**

Ramin Miri<sup>a</sup>, Hossein Sadegh pour<sup>a</sup>, Katayoun Javidnia<sup>a</sup>, Omid reza Firouzi<sup>b</sup>, Ali reza Nikkhoo<sup>c</sup>

<sup>a</sup>Department of medicinal chemistry, School of Pharmacy, Shiraz University of Medical Science

<sup>b</sup>Medicinal & Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

One third of world population is infected with latent tuberculosis (TB) infection. Based on this fact, many people are at risk to develop active TB. In 2011, tuberculosis (TB) remained the second cause of death from infectious disease worldwide. Every second one person is infected and every fifteen second someone dies with TB. Therefore, the search for new antimycobacterials is the subject of numerous recent studies.

Previous studies showed that 1, 4-dihydropyridine containing alkyl ester and diethyl carbamoyl could effectively overcome MTB. Thus, the asymmetrical analogues (6a-m) of 1, 4-dihydropyridines were synthesized by a modified Hantzsch reaction using a procedure reported by Iwanami. Reaction of acetoacetates with ammonium acetate afforded the corresponding 3-aminocrotonates. Subsequent reaction of 3-aminocrotonates with aldehydes and N, N-dimethyl acetoacetamide afforded final 1, 4-dihydropyridines (**6a-m**). The obtained compounds were purified by preparative thin layer chromatography and recrystallized from ethyl acetate-petroleum ether. They were identified by <sup>1</sup>HNMR, mass spectroscopy and IR.

These compounds have essential requirements for calcium channel blocker activity as dihydropyridine class. Therefore, these compounds were evaluated as calcium-channel antagonists by using high potassium ion concentration in guinea-pig ileal longitudinal smooth muscle. They all exhibited less calcium channel antagonist activity ( $IC_{50}=10^{-5}$  to

$10^{-6}$ ) relative to the reference drug Nifedipine ( $6.31 \times 10^{-8}$ ). Docking studies offer the binding pattern of compounds to the InhA from MTB and L-type  $\text{Ca}^{2+}$  channel. The result of pharmacological assay was used in quantitative structure-activity relationship studies utilizing multiple linear regression analysis. QSAR study indicates that the activity is related to the electronic and topological parameters. These compounds also evaluated as antimicrobial agents and they were inactive.

**Keywords:** Tuberculosis, 1, 4-dihydropyridines, calcium channel blockers, docking quantitative structure-activity relationship studie

## **Synthesis of novel asymmetrical 1, 4-dihydropyridine derivatives as anti tubercular agents and evaluation of their calcium channel blocking activity**

Ramin Miri<sup>a</sup>, Hossein Sadegh pour<sup>a</sup>, Katayoun Javidnia<sup>a</sup>, Omid reza Firouzi<sup>b</sup>, Ali reza Nikkhoo<sup>c</sup>

<sup>a</sup>Department of medicinal chemistry, School of Pharmacy, Shiraz University of Medical Science

<sup>b</sup>Medicinal & Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

One third of world population is infected with latent tuberculosis (TB) infection. Based on this fact, many people are at risk to develop active TB. In 2011, tuberculosis (TB) remained the second cause of death from infectious disease worldwide. Every second one person is infected and every fifteen second someone dies with TB. Therefore, the search for new antimycobacterials is the subject of numerous recent studies.

Previous studies showed that 1, 4-dihydropyridine containing alkyl ester and diethyl carbamoyl could effectively overcome MTB. Thus, the asymmetrical analogues (6a-m) of 1, 4-dihydropyridines were synthesized by a modified Hantzsch reaction using a procedure reported by Iwanami. Reaction of acetoacetates with ammonium acetate afforded the corresponding 3-aminocrotonates. Subsequent reaction of 3-aminocrotonates with aldehydes and N, N-dimethyl acetoacetamide afforded final 1, 4-dihydropyridines (**6a-m**). The obtained compounds were purified by preparative thin layer chromatography and recrystallized from ethyl acetate-petroleum ether. They were identified by <sup>1</sup>HNMR, mass spectroscopy and IR.

These compounds have essential requirements for calcium channel blocker activity as dihydropyridine class. Therefore, these compounds were evaluated as calcium-channel antagonists by using high potassium ion concentration in guinea-pig ileal longitudinal smooth muscle. They all exhibited less calcium channel antagonist activity ( $IC_{50}=10^{-5}$  to

$10^{-6}$ ) relative to the reference drug Nifedipine ( $6.31 \times 10^{-8}$ ). Docking studies offer the binding pattern of compounds to the InhA from MTB and L-type  $\text{Ca}^{2+}$  channel. The result of pharmacological assay was used in quantitative structure-activity relationship studies utilizing multiple linear regression analysis. QSAR study indicates that the activity is related to the electronic and topological parameters. These compounds also evaluated as antimicrobial agents and they were inactive.

**Keywords:** Tuberculosis, 1, 4-dihydropyridines, calcium channel blockers, docking quantitative structure-activity relationship studie

## **Synthesis of novel asymmetrical 1, 4-dihydropyridine derivatives as anti tubercular agents and evaluation of their calcium channel blocking activity**

Ramin Miri<sup>a</sup>, Hossein Sadegh pour<sup>a</sup>, Katayoun Javidnia<sup>a</sup>, Omid reza Firouzi<sup>b</sup>, Ali reza Nikkhoo<sup>c</sup>

<sup>a</sup>Department of medicinal chemistry, School of Pharmacy, Shiraz University of Medical Science

<sup>b</sup>Medicinal & Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

One third of world population is infected with latent tuberculosis (TB) infection. Based on this fact, many people are at risk to develop active TB. In 2011, tuberculosis (TB) remained the second cause of death from infectious disease worldwide. Every second one person is infected and every fifteen second someone dies with TB. Therefore, the search for new antimycobacterials is the subject of numerous recent studies.

Previous studies showed that 1, 4-dihydropyridine containing alkyl ester and diethyl carbamoyl could effectively overcome MTB. Thus, the asymmetrical analogues (6a-m) of 1, 4-dihydropyridines were synthesized by a modified Hantzsch reaction using a procedure reported by Iwanami. Reaction of acetoacetates with ammonium acetate afforded the corresponding 3-aminocrotonates. Subsequent reaction of 3-aminocrotonates with aldehydes and N, N-dimethyl acetoacetamide afforded final 1, 4-dihydropyridines (**6a-m**). The obtained compounds were purified by preparative thin layer chromatography and recrystallized from ethyl acetate-petroleum ether. They were identified by <sup>1</sup>HNMR, mass spectroscopy and IR.

These compounds have essential requirements for calcium channel blocker activity as dihydropyridine class. Therefore, these compounds were evaluated as calcium-channel antagonists by using high potassium ion concentration in guinea-pig ileal longitudinal smooth muscle. They all exhibited less calcium channel antagonist activity ( $IC_{50}=10^{-5}$  to

$10^{-6}$ ) relative to the reference drug Nifedipine ( $6.31 \times 10^{-8}$ ). Docking studies offer the binding pattern of compounds to the InhA from MTB and L-type  $\text{Ca}^{2+}$  channel. The result of pharmacological assay was used in quantitative structure-activity relationship studies utilizing multiple linear regression analysis. QSAR study indicates that the activity is related to the electronic and topological parameters. These compounds also evaluated as antimicrobial agents and they were inactive.

**Keywords:** Tuberculosis, 1, 4-dihydropyridines, calcium channel blockers, docking quantitative structure-activity relationship studie



## **Synthesis of novel asymmetrical 1, 4-dihydropyridine derivatives as anti tubercular agents and evaluation of their calcium channel blocking activity**

Ramin Miri<sup>a</sup>, Hossein Sadegh pour<sup>a</sup>, Katayoun Javidnia<sup>a</sup>, Omid reza Firouzi<sup>b</sup>, Ali reza Nikkhoo<sup>c</sup>

<sup>a</sup>Department of medicinal chemistry, School of Pharmacy, Shiraz University of Medical Science

<sup>b</sup>Medicinal & Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

One third of world population is infected with latent tuberculosis (TB) infection. Based on this fact, many people are at risk to develop active TB. In 2011, tuberculosis (TB) remained the second cause of death from infectious disease worldwide. Every second one person is infected and every fifteen second someone dies with TB. Therefore, the search for new antimycobacterials is the subject of numerous recent studies.

Previous studies showed that 1, 4-dihydropyridine containing alkyl ester and diethyl carbamoyl could effectively overcome MTB. Thus, the asymmetrical analogues (6a-m) of 1, 4-dihydropyridines were synthesized by a modified Hantzsch reaction using a procedure reported by Iwanami. Reaction of acetoacetates with ammonium acetate afforded the corresponding 3-aminocrotonates. Subsequent reaction of 3-aminocrotonates with aldehydes and N, N-dimethyl acetoacetamide afforded final 1, 4-dihydropyridines (**6a-m**). The obtained compounds were purified by preparative thin layer chromatography and recrystallized from ethyl acetate-petroleum ether. They were identified by <sup>1</sup>HNMR, mass spectroscopy and IR.

These compounds have essential requirements for calcium channel blocker activity as dihydropyridine class. Therefore, these compounds were evaluated as calcium-channel antagonists by using high potassium ion concentration in guinea-pig ileal longitudinal smooth muscle. They all exhibited less calcium channel antagonist activity ( $IC_{50}=10^{-5}$  to

$10^{-6}$ ) relative to the reference drug Nifedipine ( $6.31 \times 10^{-8}$ ). Docking studies offer the binding pattern of compounds to the InhA from MTB and L-type  $\text{Ca}^{2+}$  channel. The result of pharmacological assay was used in quantitative structure-activity relationship studies utilizing multiple linear regression analysis. QSAR study indicates that the activity is related to the electronic and topological parameters. These compounds also evaluated as antimicrobial agents and they were inactive.

**Keywords:** Tuberculosis, 1, 4-dihydropyridines, calcium channel blockers, docking quantitative structure-activity relationship studie